



## CLINICAL GUIDELINE

# Renal Pelvis Dilation - Detected Antenatally, Obstetrics

A guideline is intended to assist healthcare professionals in the choice of disease-specific treatments.

Clinical judgement should be exercised on the applicability of any guideline, influenced by individual patient characteristics. Clinicians should be mindful of the potential for harmful polypharmacy and increased susceptibility to adverse drug reactions in patients with multiple morbidities or frailty.

If, after discussion with the patient or carer, there are good reasons for not following a guideline, it is good practice to record these and communicate them to others involved in the care of the patient.

<b>Version Number:</b>	1
<b>Does this version include changes to clinical advice:</b>	N/A
<b>Date Approved:</b>	16 <sup>th</sup> June 2017
<b>Date of Next Review:</b>	1 <sup>st</sup> July 2022
<b>Lead Author:</b>	Chris Lilley
<b>Approval Group:</b>	Obstetrics Clinical Governance Group

### Important Note:

The Intranet version of this document is the only version that is maintained. Any printed copies should therefore be viewed as 'Uncontrolled' and as such, may not necessarily contain the latest updates and amendments.

Renal Pelvis Dilatation (Pyelectasis) – detected antenatally - GGC Alert form

**Antenatal alert to Paediatric team – refer to WoS neonatal guideline**

**Renal Pelvis Dilatation (Pyelectasis) – detected antenatally**

**Paediatric Consultant**

Local Referral centre for USS

Departmental address

Date:

Correspondence to Obstetric Secretary

**Re: maternal details**

Unit No.
CHI number
Name Sex
D.O.B.
Address
Postcode
Phone No.

**GP Details**

Name
Surgery address
Postcode

**Dear Paediatric consultant**

The above patient with an EDD of \_\_\_\_\_ has had a detailed USS at \_\_\_\_\_ weeks. Renal pelvis dilatation was evident and the measurements are as follows -

- Right \_\_\_\_\_ mm
- Left \_\_\_\_\_ mm

In addition there were the following anomalies present (please mark with an 'X') –

calyceal dilatation		oligohydramnios	
ureteral dilatation		lack of urine in bladder	
ureterocoele		bladder wall thickening	
bilateral findings		any parenchymal abnormality (detail below)	

Further details-
------------------

Any correspondence should be sent to Consultant who is the consultant in charge of the case.

**Please place a copy of this letter in the maternal notes and highlight in the paediatric alert box in the maternal notes.**

Yours sincerely

Obstetric Sonographer

Copy to:

**GP**

**Maternal Notes**

**Audit Coordinator:**

C Lilley,  
Neonatal unit, 4th Floor, Princess Royal Maternity  
16 Alexandra Parade G31 2ER