

View this article online at: [patient.info/ears-nose-throat-mouth/sore-throat-2/tonsillitis](http://patient.info/ears-nose-throat-mouth/sore-throat-2/tonsillitis)


## Tonsillitis (Symptoms and Treatment)

Tonsillitis is an inflammation of the tonsils. It is usually caused by a viral infection. Less often the cause is a bacterial infection.

Tonsillitis normally goes after a few days. Tonsillitis treatment can ease the symptoms until the infection goes. See a doctor if the symptoms are severe or don't get better quickly.

### What are tonsils?

Tonsils are made of soft gland tissue and they are part of the body's defence against infections (the immune system). You have two tonsils, one on either side at the back of the mouth.



By Blausen.com staff (2014). "Medical gallery of Blausen Medical 2014". WikiJournal of Medicine 1 (2). DOI:10.15347/wjm/2014.010. ISSN 2002-4436., CC BY 3.0, via Wikimedia Commons

### What are the symptoms of tonsillitis?

A sore throat is the most common of all tonsillitis symptoms. You may also cough or sneeze, have high temperature (fever), headache, feel sick (nausea), feel tired, find swallowing painful, and have swollen lymph nodes in the neck. The tonsils may swell and become red. Pus may appear as white spots on the tonsils. Symptoms typically get worse over 2-3 days and then gradually go, usually within a week. The picture below shows inflamed tonsils.



### What causes tonsillitis?

Tonsillitis is sometimes caused by a bacterial infection like strep throat (group A Streptococcus), but usually by a cold or flu. Swollen tonsils may also sometimes cause obstructive sleep apnoea (OSA).

### What is glandular fever?

Glandular fever (infectious mononucleosis) is caused by a virus (the Epstein-Barr virus). It tends to cause a severe bout of tonsillitis as well as other symptoms. [See the separate leaflet called Glandular Fever \(Infectious Mononucleosis\) for more details.](#)

## Is tonsillitis contagious?

Contagious means you can pass the infection to someone else. Tonsillitis is contagious so infection may indeed spread by close physical contact with other people, or by droplets in the air, caused by sneezing or coughing.

As with [coughs](#), [colds](#), [flu](#) and other similar infections, there is a chance that you can pass on the virus or bacterium that has caused tonsillitis.

## When should you contact a doctor about tonsillitis?

Seek medical advice if symptoms of a sore throat cause severe symptoms, or if they do not ease within 3-4 days. In particular, you should seek urgent medical attention if you develop:

- Difficulty in breathing.
- Difficulty swallowing saliva.
- Difficulty opening your mouth.
- Severe pain.
- A persistent high temperature.
- A severe illness, especially when symptoms are mainly on one side of the throat.

## Tonsillitis treatments

Mild tonsillitis often doesn't need any treatment. However, it is important to drink plenty of water. Paracetamol or ibuprofen will help to ease pain, headache and high temperature. Gargles, lozenges and sprays may help to soothe a sore throat but they do not shorten the illness.

Most throat and tonsil infections are caused by viruses but some are caused by bacteria. Antibiotics kill bacteria but do not kill viruses. Even if tonsillitis is caused by a bacterium, treatment with an antibiotic does not make much difference in most cases.

If you have repeated bouts of tonsillitis you may consider having your tonsils removed (tonsillectomy).

Having no treatment at all is one option. Many tonsil infections are mild and soon get better on their own.

Treatment options include.

- **Have plenty to drink.** It is tempting not to drink very much if it is painful to swallow. You may become mildly lacking in fluid in the body (mildly dehydrated) if you don't drink much, particularly if you also have a high temperature (fever). Some lack of fluid in the body can make headaches and tiredness much worse.
- **Paracetamol or ibuprofen** eases pain, headache and fever. To keep symptoms to a minimum it is best to take a dose at regular intervals as recommended on the packet of medication rather than now and then. For example, take [paracetamol](#) four times a day until symptoms ease. Although either paracetamol or [ibuprofen](#) will usually help, there is some evidence to suggest that ibuprofen may be more effective than paracetamol at easing symptoms in adults. Paracetamol is usually the preferred first-line option for children, but ibuprofen can be used as an alternative. **Note:** some people with certain conditions may not be able to take ibuprofen. So, always read the packet label.
- **Other gargles, lozenges and sprays** that you can buy at pharmacies may help to soothe a sore throat. However, they do not shorten the illness.

## Do you need antibiotics for tonsillitis?

Usually not. Most throat and tonsil infections are caused by germs called viruses, although some are caused by germs called bacteria. Without tests, it is usually not possible to tell if it is a viral or bacterial tonsillitis. Antibiotics kill bacteria, but do not kill viruses.

Even if a bacterium is the cause of a tonsil or throat infection, an antibiotic does not make much difference in most cases. Your body defences (immune system) usually clear these infections within a few days whether caused by a virus or a bacterium. Also, antibiotics can sometimes cause side-effects such as diarrhoea, rash and stomach upsets.

So, most doctors do not prescribe antibiotics for most cases of tonsillitis or sore throat.

An antibiotic may be advised in certain situations. For example:

- If the infection is severe.
- If it is not easing after a few days.
- If your immune system is not working properly (for example, if you have had your [spleen removed](#), if you are taking [chemotherapy](#), etc).

## Do I need my tonsils removing?

If you have repeated (recurring) tonsillitis you may wonder about having your tonsils removed. Guidelines suggest removing the tonsils may be an option (tonsillectomy) if you:

- Have had seven or more episodes of tonsillitis in the preceding year; **or**
- Five or more such episodes in each of the preceding two years; **or**
- Three or more such episodes in each of the preceding three years.
- **And ...**
- The bouts of tonsillitis affect normal functioning. For example, they are severe enough to make you need time off from work or for young children taking time off from school.

The tonsils and adenoids may be removed at the same time. The adenoids are also part of the body's defence against infections (the immune system). Adenoids hang from the upper part of the back of the nasal cavity. [See the separate leaflet called What do tonsils do? for more details.](#)

Although full-blown episodes of tonsillitis are prevented after tonsillectomy, other throat infections are not prevented. However, the overall number and severity of throat infections may be reduced. Also, the risk of developing quinsy is reduced. Many people say they generally feel better in themselves after having their tonsils removed if they previously had frequent episodes of chronic tonsillitis.

Tonsillectomy is usually a straightforward minor operation. But, as with all operations, there is a risk. For example, there is a small risk of life-threatening severe bleeding from the throat during or just after the operation.

## What is the outlook for tonsillitis?

In nearly all cases, a tonsillitis or sore throat clears away without leaving any problems. However, occasionally tonsillitis may progress to cause a complication. Also, a sore throat or tonsillitis is sometimes due to an unusual, but more serious, illness.

Sometimes the infection can spread from the tonsils to other nearby tissues. For example, to cause an [ear infection](#), [sinus infection](#) or [chest infection](#).

## What is quinsy?

Quinsy is also known as peritonsillar abscess. Quinsy is an uncommon condition where a collection of pus (abscess) develops next to a tonsil. This is caused by a bacterial infection.

- It usually develops just on one side. It may follow a tonsillitis or develop without tonsillitis.
- The tonsil on the affected side may be swollen or look normal, but is pushed towards the middle by the abscess next to the tonsil. Quinsy is very painful and can make you feel very unwell.
- It is treated with antibiotics, but also the pus often needs to be drained with a small operation.

## Further reading & references

- [Management of sore throat and indications for tonsillectomy](#); Scottish Intercollegiate Guidelines Network - SIGN (April 2010)
- [Sore throat \(acute\): antimicrobial prescribing](#); NICE Guideline (January 2018)
- [Burton MJ, Glasziou PP, Chong LY, et al](#); Tonsillectomy or adenotonsillectomy versus non-surgical treatment for chronic/recurrent acute tonsillitis. *Cochrane Database Syst Rev.* 2014 Nov 19;(11):CD001802. doi: 10.1002/14651858.CD001802.pub3.
- [Spinks A, Glasziou PP, Del Mar CB](#); Antibiotics for treatment of sore throat in children and adults. *Cochrane Database Syst Rev.* 2021 Dec 9 ;11:CD000023.
- [Tonsillectomy - Commissioning Guide](#); ENT UK and Royal College of Surgeons, 2021

**Disclaimer:** This article is for information only and should not be used for the diagnosis or treatment of medical conditions. Egton Medical Information Systems Limited has used all reasonable care in compiling the information but makes no warranty as to its accuracy. Consult a doctor or other healthcare professional for diagnosis and treatment of medical conditions. For details see our [conditions](#).

Author: Dr Laurence Knott	Peer Reviewer: Dr Hayley Willacy	
Document ID: 4403 (v47)	Last Checked: 21/07/2021	Next Review: 20/07/2024

View this article online at: [patient.info/ears-nose-throat-mouth/sore-throat-2/tonsillitis](http://patient.info/ears-nose-throat-mouth/sore-throat-2/tonsillitis)

Discuss Tonsillitis (Symptoms and Treatment) and find more trusted resources at [Patient](#).



**Book appointments,  
order repeat prescriptions and  
view your medical record online**

To find out more visit  
[www.patientaccess.com](http://www.patientaccess.com)  
or download the app

