


4

Patient Information on

# Managing Breathlessness

**Leaflet 4:** Positions to ease breathlessness



**The information given below is designed to help you manage your stable long term breathlessness. If your breathing is getting worse or you are experiencing breathlessness as a new feeling, it is important to seek medical advice from your GP.**

## When to use

These positions may help you to ease your breathlessness at rest and after activity. The most commonly used positions are shown, however you may find other positions of ease which is not pictured. Experiment with and modify different positions until you find what works best for you.

## How positioning helps to ease breathlessness

The positions in this leaflet place the arms so the breathing accessory muscles are in a better position to help with breathing. Leaning forward may also improve the movement of your diaphragm, the main muscle of breathing. For more information on muscles of breathing see Leaflet 3

## Positions to ease breathlessness following activity

The following positions may help to ease breathlessness after you have been active. With all positions, try to relax the hands, wrists, shoulders, neck and jaw as much as possible. Experiment with your arm position. Does your breathing feel easier with your hands behind your head or back?



There is no one position which suits everyone, nor one position which people with a particular condition prefer. Experiment with different positions and find what works best for you.

## Use of walking aids

Some people may notice that they are able to walk further and feel less breathless when pushing a supermarket trolley around the shop. Pushing the trolley enables you to maintain a 'forward lean position of ease' while walking.

In a similar way, using a walking frame may enable you to walk further and feel less breathless, as well as providing a portable leaning post when you wish to stop to rest. A walking stick can also provide a portable leaning post.



**Gutter frame**



**Rotator frame**



**Walking stick**

When walking outdoors, people often prefer to use a 3 or 4 wheeled walker. Some walkers have a seat so you can sit to rest, others may have a handy storage compartment.

**This leaflet has been produced by the Cambridge Breathlessness Intervention Service [www.cuh.org.uk/bis](http://www.cuh.org.uk/bis) and is reproduced here in NHS Highland with their kind permission. Other leaflets in this series will go through these techniques in a bit more detail.**

## Document history

<b>Authors</b>	Breathlessness Intervention Service
<b>Pharmacist</b>	n/a
<b>Department</b>	Cambridge University Hospitals NHS Foundation Trust, Hills Road, Cambridge, CB2 0QQ <a href="http://www.cuh.org.uk">www.cuh.org.uk</a>
<b>Contact number</b>	01223 274404
<b>Publish/review date</b>	September 2016/September 2019
<b>File name</b>	Leaflet_4_positions_to_ease_breathlessness.doc
<b>Version number/Ref</b>	1/100080