
Wound Cleansing Pathway



Cleansing solutions should be at room temperature.

*Potable water is water pure enough to be consumed or used with low risk of immediate or long term harm e.g. tap water. For private water supplies where quality cannot be guaranteed, we would suggest using NaCl 0.9%.

**Biofilm disrupting cleansing solution has specific activity to breakdown or inhibit biofilm growth e.g. PHMB Polyhexanide and betaine is a polymer used as a wound cleanser.



References:

Brown, A. (2018) When is wound cleansing necessary and what solution should be used? Nursing Times, Vol. 114 (9), pp. 40 -43.

Flanagan, M. (2013) Principles of wound management. In: Flanagan M (ed) Wound Healing and Skin Integrity: Principles and Practice. Chichister: Wiley-Blackwell.

Collier, M. and Hofer, P. (2017) Taking wound cleansing seriously to minimise risk. Wounds UK Vol 13 (1) pp 58- 64.

Hűbner, N.O., Kramer, A. (2010) Review on the efficacy, safety and clinical applications of polihexanide, a modern wound antiseptic. Skin Pharmacology and Physiology, Vol 23, Suppl, pp17-27.

Kaehn, K., Eberlein, T. (2009) In-vitro test for comparing the efficacy of wound rinsing solutions. British Journal of Nursing (Tissue Viability Supplement) Vol 18, No. 11 (S4 – S9). Wolcott, R. Fletcher, J (2014) The Role of wound cleansing in the management of wounds. Wounds International, Vol. 5 (3), pp. 25-31.